

項目毎の最高グレード		グレード				発現率(%)		
		1	2	3	4	≥G1	≥G2	≥G3
臨床値	血色素減少	14	24	10	2	90.9	65.5	21.8
	赤血球減少	12	14	16	1	78.2	56.4	30.9
	ヘマトクリット値減少	9	16	11	1	67.3	50.9	21.8
	白血球減少	8	18	2	1	52.7	38.2	5.5
	好中球減少	7	13	13	3	65.5	52.7	29.1
	血小板減少	28	4	1		60.0	9.1	1.8
	リンパ球減少		8	3		20.0	20.0	5.5
	単球増多	1				1.8		
	AST上昇	7		1		14.5	1.8	1.8
	ALT上昇	6	1	1		14.5	3.6	1.8
	ALP上昇	5	1			10.9	1.8	
	LDH上昇	4				7.3		
	総蛋白低下	25				45.5		
	アルブミン低下	16	6			40.0	10.9	
	血清ビリルビン上昇	9	7			29.1	12.7	
	BUN上昇	12				21.8		
	血中クレアチニン上昇	9	1	1		20.0	3.6	1.8
	血清ナトリウム上昇	2				3.6		
	血清カリウム上昇	7	1			14.5	1.8	
	血清クロール上昇	2				3.6		
	血清ナトリウム低下	12		4		29.1	7.3	7.3
	血清カリウム低下	3		2		9.1	3.6	3.6
	血清クロール低下	11	1	1		23.6	3.6	1.8
	血清カルシウム低下	1				1.8		
	尿糖陽性	4	2			10.9	3.6	
	蛋白尿	11	1			21.8	1.8	
	ウロビリノーゲン陽性	2				3.6		
	血糖値上昇			1		1.8	1.8	1.8
	CCr低下	4	1	1		10.9	3.6	1.8

グレード：NCI-CTC ver.2.0
副作用用語はJ-ART により分類した。

項目毎の最高グレード		グレード				発現率(%)		
		1	2	3	4	≥G1	≥G2	≥G3
臨床症状	食欲不振	21	15	7		78.2	40.0	12.7
	悪 心	22	8	6		65.5	25.5	10.9
	嘔 吐	14	3	4		38.2	12.7	7.3
	下 痢	12	3	4		34.5	12.7	7.3
	腹 痛	2	1			5.5	1.8	
	腹部不快感	1				1.8		
	心窩部痛		1			1.8	1.8	
	胸やけ	1				1.8		
	しゃっくり	1	3			7.3	5.5	
	便 秘	2	1	2		9.1	5.5	3.6
	胃潰瘍			1		1.8	1.8	1.8
	口内炎	12	2			25.5	3.6	
	口角炎	1				1.8		
	発 疹	3	2			9.1	3.6	
	落 屑	3	1			7.3	1.8	
	顔面潮紅	1				1.8		
	瘙 痒	3				5.5		
	脱 毛	5				9.1		
	色素沈着	13				23.6		
	爪の変色	1	1			3.6	1.8	
	爪の変形		1			1.8	1.8	
	爪の隆起	1				1.8		
	頭 痛	1				1.8		
	頭重感	1				1.8		
	不 眠	1				1.8		
	失神発作			2		3.6	3.6	3.6
	全身倦怠感	5	1			10.9	1.8	
	疲 労	17	8			45.5	14.5	
	発 熱	4	1			9.1	1.8	
	体重減少	7	2			16.4	3.6	
	血圧上昇			1		1.8	1.8	1.8
	下肢浮腫	1				1.8		
	尿 閉			1		1.8	1.8	1.8
	鼻出血	1				1.8		
	気道感染	1				1.8		
	流 涙	3				5.5		
	結膜炎	1				1.8		
	角膜炎			1		1.8	1.8	1.8
	鼻 汁	1				1.8		
	難 聴	1				1.8		
	耳 鳴	1				1.8		
	味覚異常	1				1.8		
	咽頭炎	2				3.6		
	胸 水		1			1.8	1.8	

(2004年12月非小細胞肺癌効能追加時までの集計
大鵬薬品工業株式会社)